

LU-DECOMPOSITION OF A NONCOMMUTATIVE LINEAR SYSTEM AND JACOBI POLYNOMIALS

OSCAR BREGA AND LEANDRO CAGLIERO

ABSTRACT. In this paper we obtain the LU-decomposition of a noncommutative linear system of equations that, in the rank one case, characterizes the image of the Lepowsky homomorphism $U(\mathfrak{g})^K \rightarrow U(\mathfrak{k})^M \otimes U(\mathfrak{a})$. This LU-decomposition can be transformed into very simple matrix identities, where the entries of the matrices involved belong to a special class of Jacobi polynomials. In particular, each entry of the L part of the original system is expressed in terms of a single ultraspherical Jacobi polynomial. In turns, these matrix identities yield a biorthogonality relation between the ultraspherical Jacobi polynomials.

1. INTRODUCTION

1.1. The noncommutative linear system. Let k be a field of characteristic zero, let $\mathcal{A}$ be an associative, not necessarily commutative, k -algebra with unit and let $\mathcal{M}$ be a unital $\mathcal{A}$ -bimodule.

This paper is devoted to perform the gaussian elimination process on the following non commutative homogeneous system of infinitely many linear equations and infinite unknowns

$$(1.1) \quad \begin{array}{ccccccc} Ep_0 & + & Ep_1 & + & Ep_2 & \dots & = & p_0E & - & p_1E & + & p_2E & \dots \\ E^2p_0 & + & 2E^2p_1 & + & 2^2E^2p_2 & \dots & = & p_0E^2 & - & 2p_1E^2 & + & 2^2p_2E^2 & \dots \\ E^3p_0 & + & 3E^3p_1 & + & 3^2E^3p_2 & \dots & = & p_0E^3 & - & 3p_1E^3 & + & 3^2p_2E^3 & \dots \\ \vdots & & \vdots & & \vdots & & & \vdots & & \vdots & & \vdots & \end{array} ,$$

where E is a given element of $\mathcal{A}$ and $p_0, p_1, p_2, \dots$ belong to $\mathcal{M}$ and are the unknowns.

It is clear that $(p_0, p_1, \dots, p_d, 0, \dots)$ is a solution of the system (1.1) if and only if the polynomial $p = p_0 + p_1t + p_2t^2 + \dots + p_dt^d$, belonging to $\mathcal{M}[t] := \mathcal{M} \otimes k[t]$, satisfies the equations

$$(1.2) \quad E^n p(n) = p(-n) E^n \quad \text{for all } n \in \mathbb{N}.$$

This system can not be expressed as a single matrix equation $AX = 0$ with A a matrix with coefficients in $\mathcal{A}$. In fact, non commutative systems of (homogeneous) linear equations can seldom be expressed as a single matrix equation with coefficients in $\mathcal{A}$. Even when it is possible, only in exceptional cases a gaussian elimination process can be performed in a satisfactory way. For instance, if $\dim_k \mathcal{A} < \infty$ and $\dim_k \mathcal{M} < \infty$, by using bases of $\mathcal{A}$ and $\mathcal{M}$, one can express every finite system of linear equations as a single matrix equation $AX = B$ with coefficients in k . However the size of A might be so

big that this approach does not help very much. Some papers dealing with this subject are [C], [CS], [Or] or [GGRW].

On the other hand, by using the left and right regular actions $L, R : \mathcal{A} \rightarrow \text{End}_k(\mathcal{M})$ of $\mathcal{A}$ on $\mathcal{M}$, it is indeed possible to express every non commutative system of linear equations as a single matrix equation but with coefficients in $\text{End}_k(\mathcal{M})$. In our case, the system (1.1) can be expressed as single matrix

$$\text{equation } M_0 X = 0, \text{ where } X = \begin{pmatrix} p_0 \\ p_1 \\ p_2 \\ \vdots \end{pmatrix} \text{ and}$$

$$M_0 = \begin{pmatrix} L_E - R_E & L_E + R_E & L_E - R_E & L_E + R_E & \dots \\ L_E^2 - R_E^2 & 2L_E^2 + 2R_E^2 & 4L_E^2 - 4R_E^2 & 8L_E^2 + 8R_E^2 & \dots \\ L_E^3 - R_E^3 & 3L_E^3 + 3R_E^2 & 9L_E^3 - 9R_E^2 & 27L_E^3 + 27R_E^2 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

In order to have an insight into the LU-decomposition of this system assume for a moment that the unknown polynomial has degree 2. In this case it is easy to see that

$$\begin{pmatrix} L_E - R_E & L_E + R_E & L_E - R_E \\ L_E^2 - R_E^2 & 2L_E^2 + 2R_E^2 & 4L_E^2 - 4R_E^2 \\ L_E^3 - R_E^3 & 3L_E^3 + 3R_E^2 & 9L_E^3 - 9R_E^2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ L_E + R_E & 1 & 0 \\ L_E^2 + R_E L_E + R_E^2 & 2L_E + 2R_E & 1 \end{pmatrix} \begin{pmatrix} L_E - R_E & L_E + R_E & L_E - R_E \\ 0 & (L_E - R_E)^2 & 3L_E^2 - 3R_E^2 \\ 0 & 0 & 2(L_E - R_E)^3 \end{pmatrix}.$$

is the LU-decomposition of the system (1.1). The goal of this paper is to obtain a transparent expression for the LU-decomposition of the following more general system

$$(1.3) \quad E^n p(H + n) = p(H - n) E^n \quad \text{for all } n \in \mathbb{N},$$

for some $H \in \mathcal{A}$.

1.2. Relevance of the system (1.3). The interest on the system (1.3) comes from the fact that their solution set is closely related to invariant spaces under group actions. This is clear in the particular case in which $H = 0$ and E is the identity of $\mathcal{A}$, since the solutions of (1.3) are the even polynomials. It is not difficult to see that in other more general situations there exists a group G_E associated to E , acting on $\mathcal{M} \otimes_k k[t]$, such that

$$\{p \in \mathcal{M} \otimes_k k[t] : E^n p(n) = p(-n) E^n \text{ for all } n \in \mathbb{N}\} = (\mathcal{M} \otimes_k k[t])^{G_E}.$$

A second, and more relevant, example of the relationship between the system (1.3) and invariants of groups appears in a problem from representation theory of Lie groups. Let G_o be a connected, noncompact, real semisimple Lie group with finite center, let $\mathfrak{g}$ be the complexification of the Lie algebra of G_o and let K_o denote a maximal compact subgroup of G_o . By the fundamental work of Harish-Chandra it is known that many deep questions concerning the infinite dimensional representation theory of G_o reduce to questions about the structure of the algebra $U(\mathfrak{g})^{K_o}$, the K_o -invariants in the universal enveloping algebra of $\mathfrak{g}$. In [T], Tirao proved that the elements

of the image of $U(\mathfrak{g})^{K_\circ}$ by the Lepowsky homomorphism (see [L]) satisfy a system of linear equations completely analogous to system (1.3). Moreover, for rank one classical Lie groups it is proved in [BCT] that this image coincides with the solution set of the system introduced by Tirao. In order to prove this result, the LU-decomposition obtained in this paper constitute a very useful tool. These connections with groups invariants are not treated in this paper.

1.3. Main results. In §2 we find a periodic sequence of elementary row operations that, when applied to the system (1.3), transforms it into a triangular system that involves the adjoint action of E and some combinatoric sums of discrete derivatives of the polynomial p . The main result of this section is Theorem 2.2. As a corollary we obtain that if $p \in \mathcal{M}[t]$ satisfies

$$E^n p(H + n) = p(H - n) E^n \quad \text{for } 0 \leq n \leq \deg(p),$$

then p satisfies (1.3).

Note that M_0 is an infinite matrix with coefficients in $\text{End}_k(\mathcal{M})$ and it is closely related to the matrix

$$\tilde{M}_0 = \begin{pmatrix} x-1 & x+1 & x-1 & x+1 & \cdot \\ x^2-1 & 2x^2+2 & 4x^2-4 & 8x^2+8 & \cdot \\ x^3-1 & 3x^3+3 & 9x^3-9 & 27x^3+27 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix} \in k[x].$$

Taking into account the relationship between M_0 and $\tilde{M}_0$, the results obtained in Theorem 2.2 can be translated in terms of polynomials in one variable with coefficients in k . The combinatorial aspects of Theorem 2.2 are transformed into very simple matrix identities, where the entries of the matrices involved are a special class of Jacobi polynomials. In turns, these matrix identities yield biorthogonality relations between the ultraspherical polynomials. It has recently appeared in the literature some other matrix identities (in particular LU decompositions) involving Jacobi polynomials that translate sophisticated polynomials identities into very simple matrix identities (see for instance [KO]).

These results are described in what follows.

Let $P_n^{\alpha,\beta}$ denote the Jacobi polynomial of degree n associated to the numbers α and β . Usually α and β are complex numbers but, since we are working in the context of an arbitrary field of characteristic zero, they are assumed to be rational. Recall that the polynomials corresponding to parameters $\alpha = \beta$ are known as the *ultraspherical* or *Gegenbauer's* polynomials. Let

$$p_n^{\alpha,\beta}(x) = (x-1)^n P_n^{\alpha,\beta}\left(\frac{x+1}{x-1}\right).$$

It is clear that $p_n^{\alpha,\beta}(x)$ is again a polynomial and it can be expressed in terms of the hypergeometric function of Gauss as follows (see (4.22.1) in [Sz]),

$$p_n^{\alpha,\beta}(x) = \frac{(n+\alpha+\beta+1)\dots(2n+\alpha+\beta)}{n!} {}_2F_1(-n, -n-\alpha; -2n-\alpha-\beta; x-1).$$

Our results in terms of $\tilde{M}_0$ are the following:

(1) The LU-decomposition of $\tilde{M}_0$ is

$$\tilde{M}_0 = \tilde{L}_0 \tilde{U}_0 \begin{pmatrix} 1 & & & & & \\ & 1 & & & & \\ & & 2 & & & \\ & & & 6 & & \\ & & & & 24 & \\ & & & & & \end{pmatrix} \begin{pmatrix} 1 & -1 & 1 & -1 & 1 & \cdot \\ & 1 & -3 & 7 & -15 & \cdot \\ & & 1 & -6 & 25 & \cdot \\ & & & 1 & -10 & \cdot \\ & & & & 1 & \cdot \\ & & & & & \cdot \end{pmatrix},$$

where

$$(\tilde{L}_0)_{ij} = (-1)^{i-j} p_{i-j}^{-i,-i}(x), \quad i \geq j;$$

$$(\tilde{U}_0)_{ij} = (-1)^{i-j} \frac{j}{i} (x-1)^{2i-j} p_{j-i}^{-j,-1}(x), \quad i \leq j;$$

and the explicit numerical matrices are respectively formed by the factorial and (up to the minus signs) Stirling numbers of the second kind.

(2) The inverse of the matrix $\tilde{L}_0$ is

$$(\tilde{L}_0^{-1})_{ij} = (-1)^{i-j} \frac{j}{i} p_{i-j}^{j,j}(x), \quad i \geq j.$$

This yields the following “discrete orthogonality” relationship that involves once many of the ultraspherical Jacobi polynomials with integer parameters

$$\begin{pmatrix} \frac{1}{1} P_0^{1,1} & & & & \\ \frac{1}{2} P_1^{1,1} & \frac{2}{2} P_0^{2,2} & & & \\ \frac{1}{3} P_2^{1,1} & \frac{2}{3} P_1^{2,2} & \frac{3}{3} P_0^{3,3} & & \\ \frac{1}{4} P_3^{1,1} & \frac{2}{4} P_2^{2,2} & \frac{3}{4} P_1^{3,3} & \frac{1}{1} P_0^{4,4} & \\ & & & & \end{pmatrix} \begin{pmatrix} P_0^{-1,-1} & & & & \\ P_1^{-2,-2} & P_0^{-2,-2} & & & \\ P_2^{-3,-3} & P_1^{-3,-3} & P_0^{-3,-3} & & \\ P_3^{-4,-4} & P_2^{-4,-4} & P_1^{-4,-4} & P_0^{-4,-4} & \\ & & & & \end{pmatrix} = \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & \end{pmatrix}.$$

(3) We mentioned above that we performed to the system (1.3) a gaussian elimination process in which we periodically apply the same sequence of elementary row operations. This periodicity is translated in the following terms. Given an infinite matrix

$$T = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdot \\ a_{21} & a_{22} & a_{23} & \cdot \\ a_{31} & a_{32} & a_{33} & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix},$$

let $s(T)$ be the *shifted matrix* of T , that is $s(T)$ is the diagonal blocked matrix $\begin{pmatrix} 1 & \\ & T \end{pmatrix}$ formed by the identity matrix of size 1×1 and T . We say that an infinite matrix A admits a *periodic gaussian elimination process* if there exist an lower triangular matrix with 1's in the diagonal T_0 such that the sequence of matrices

$$A, \quad T_0 A, \quad s(T_0) T_0 A, \quad s^2(T_0) s(T_0) T_0 A, \quad \dots$$

converges¹ to an upper triangular matrix. In this case T_0 is called a *fundamental sequence* of the Gaussian elimination process of A . It is not difficult to see that an infinite matrix A admits a periodic gaussian elimination process if and only if A admits an LU-decomposition. In

¹The concept of convergence is used in the discrete sense, that is a sequence of matrices B_k converges to B if for every i, j there exists k_0 such that $(B_k)_{ij} = B_{ij}$ for all $k \geq k_0$.

fact, if $A = LU$ then the fundamental period is $T_0 = s(L)L^{-1}$ and, conversely, if T_0 is the fundamental period of A then

$$L^{-1} = \cdots s^2(T_0)s(T_0)T_0 \quad \text{and} \quad L = T_0^{-1}s(T_0^{-1})s^2(T_0^{-1})\cdots.$$

It is worth mentioning at this point that the Vandermonde matrix $V_{ij} = i^{j-1}$ admits a periodic gaussian elimination process and its fundamental sequence is $\begin{pmatrix} 1 & & & \\ -1 & 1 & & \\ & -1 & 1 & \\ & & -1 & 1 \end{pmatrix}$ and that the recursive definition of the Pascal numbers is equivalent to the identity

$$\begin{pmatrix} 1 & & & \\ 1 & 1 & & \\ 1 & 2 & 1 & \\ 1 & 3 & 3 & 1 \end{pmatrix} = \cdots s^2(T_0)s(T_0)T_0, \quad \text{with } T_0 = \begin{pmatrix} 1 & & & \\ 1 & 1 & & \\ & 1 & 1 & \\ & & 1 & 1 \end{pmatrix}.$$

It turns out that $\tilde{M}_0$ admits a periodic gaussian elimination process and we show that its fundamental sequence is

$$T_0 = \begin{pmatrix} 1 & & & \\ -x-1 & 1 & & \\ x & -x-1 & 1 & \\ & x & -x-1 & 1 \end{pmatrix}.$$

It is not difficult to see that the inverse of T_0 is the matrix formed by the cyclotomic polynomials

$$T_0^{-1} = \begin{pmatrix} 1 & & & \\ 1+x & 1 & & \\ 1+x+x^2 & 1+x & 1 & \\ 1+x+x^2+x^3 & 1+x+x^2 & 1+x & 1 \end{pmatrix},$$

and therefore the ultraspherical polynomials can be expressed as

$$\begin{pmatrix} \frac{1}{1}p_0^{1,1} & & & \\ \frac{1}{2}p_1^{1,1} & \frac{2}{2}p_0^{2,2} & & \\ \frac{1}{3}p_2^{1,1} & \frac{2}{3}p_1^{2,2} & \frac{3}{3}p_0^{3,3} & \\ \frac{1}{4}p_3^{1,1} & \frac{2}{4}p_2^{2,2} & \frac{3}{4}p_1^{3,3} & \frac{1}{1}p_0^{4,4} \end{pmatrix} \\ = \cdots \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & -x-1 & 1 & \end{pmatrix} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & -x-1 & 1 & \\ & x & -x-1 & 1 \end{pmatrix} \begin{pmatrix} 1 & & & \\ -x-1 & 1 & & \\ x & -x-1 & 1 & \\ & x & -x-1 & 1 \end{pmatrix}$$

and

$$\begin{pmatrix} p_0^{-1,-1} & & & \\ p_1^{-2,-2} & p_0^{-2,-2} & & \\ p_2^{-3,-3} & p_1^{-3,-3} & p_0^{-3,-3} & \\ p_3^{-4,-4} & p_2^{-4,-4} & p_1^{-4,-4} & p_0^{-4,-4} \end{pmatrix} \\ = \begin{pmatrix} 1 & & & \\ 1+x & 1 & & \\ 1+x+x^2 & 1+x & 1 & \\ 1+x+x^2+x^3 & 1+x+x^2 & 1+x & 1 \end{pmatrix} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & 1+x & 1 & \\ & 1+x+x^2 & 1+x & 1 \end{pmatrix} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1+x \end{pmatrix} \cdots$$

We notice that in these two identities the polynomials involved are $p_n^{\alpha,\alpha}$ instead of $P_n^{\alpha,\alpha}$.

2. THE GAUSSIAN ELIMINATION FOR THE SYSTEM (1.3)

Let k be a field of characteristic zero, let $\mathcal{A}$ be an associative, not necessarily commutative, k -algebra with unit and let $\mathcal{M}$ be a unital $\mathcal{A}$ -bimodule. Given an element $r \in \mathcal{A}$, let $L_r, R_r \in \text{End}_k(\mathcal{M})$ respectively denote the left and right actions by r , and let $\text{ad}_r = L_r - R_r$ be the adjoint action of r in $\mathcal{M}$.

Recall that $\mathcal{M}[t] = \mathcal{M} \otimes_k k[t]$, and that given an element $r \in \mathcal{A}$ one has the evaluation map $\mathcal{M}[t] \rightarrow \mathcal{M}$ defined using the right action of $\mathcal{A}$ on $\mathcal{M}$ by $a \otimes p(t) \mapsto ap(r)$. If $r \in \mathcal{A}$ and $p \in \mathcal{M}[t]$ then the evaluation of p in r is given by $p(r) = p_0 + p_1 r + \cdots + p_n r^n$.

Given two arbitrary elements E and H in $\mathcal{A}$ we are interested in the set of polynomials $p \in \mathcal{M}[t]$ that satisfy the following system of linear equations (1.3), that is polynomials $p \in \mathcal{M}[t]$ satisfying

$$E^n p(H + n) = p(H - n) E^n \quad \text{for all } n \in \mathbb{N}.$$

In this section, we assume that E is an eigenvector of ad_H in $\mathcal{A}$ with eigenvalue c and we shall perform a gaussian elimination process to obtain, in Theorem 2.2, a triangular linear system in $\mathcal{M}[t]$ equivalent to the system (1.3).

2.1. A first look at the system. We first discuss the linear system (1.3) for polynomials of degree zero and one.

Degree zero: Let $p = p_0 \in \mathcal{M}[t]$ be a constant polynomial. Observe that p satisfies the first equation of the system if and only if p_0 commutes with E , and therefore p satisfies all the other equations. Thus p satisfies the system (1.3) if and only if $p_0 \in \ker \text{ad}_E$.

Degree one: A linear polynomial $p = p_0 + p_1 t \in \mathcal{M}[t]$ satisfies the system (1.3) if and only if the vector (p_0, p_1) is a solution of the following system,

$$(2.1) \quad E p_0 + E p_1 (H+1) = p_0 E + p_1 (H-1) E$$

$$(2.2) \quad E^2 p_0 + E^2 p_1 (H+2) = p_0 E^2 + p_1 (H-2) E^2$$

$$(2.3) \quad E^3 p_0 + E^3 p_1 (H+3) = p_0 E^3 + p_1 (H-3) E^3$$

$$\vdots \qquad \qquad \qquad \vdots$$

It is not difficult to see that equation (2.3) can be obtained from the first two as follows,

$$2E \times (2.2) - E^2 \times (2.1) - E \times (2.1) \times E - (2.1) \times E^2 + (2.2) \times E.$$

In fact, we shall see in Corollary 2.3 that for all $n \geq 3$, the n th equation can be obtained from equations (2.1) and (2.2). Then, in this case, the system (1.3) is equivalent to the 2×2 system formed by equations (2.1) and (2.2). This 2×2 system is represented by the matrix

$$M = \begin{pmatrix} L_E - R_E & L_E R_{H+1} - R_E R_{H-1} \\ L_E^2 - R_E^2 & L_E^2 R_{H+2} - R_E^2 R_{H-2} \end{pmatrix},$$

with coefficients in $\text{End}_k(\mathcal{M})$. Now, since

$$\begin{pmatrix} 1 & 0 \\ -(L_E + R_E) & 1 \end{pmatrix} \begin{pmatrix} L_E - R_E & L_E R_{H+1} - R_E R_{H-1} \\ L_E^2 - R_E^2 & L_E^2 R_{H+2} - R_E^2 R_{H-2} \end{pmatrix} = \begin{pmatrix} L_E - R_E & L_E R_{H+1} - R_E R_{H-1} \\ 0 & (L_E - R_E)^2 \end{pmatrix},$$

we obtain that equations (2.1) and (2.2), are equivalent to the triangular system $U \begin{pmatrix} b_0 \\ b_1 \end{pmatrix} = 0$, with

$$U = \begin{pmatrix} L_E - R_E & L_E R_{H+1} - R_E R_{H-1} \\ 0 & (L_E - R_E)^2 \end{pmatrix} = \begin{pmatrix} L_E - R_E & L_E + R_E \\ 0 & (L_E - R_E)^2 \end{pmatrix} \begin{pmatrix} 1 & R_H \\ 0 & 1 \end{pmatrix}.$$

In Theorem 2.2 we shall extend this triangularization process to polynomials of arbitrary degree. In particular, we shall obtain in Corollary 2.3 that for polynomials of degree N the whole system (1.3) is equivalent to the first $N + 1$ equations of the system (1.3).

2.2. Some additional notation. For $h \in \mathbf{k}$, $h \neq 0$, we define the h -discrete derivative $\partial_h : \mathcal{M}[t] \rightarrow \mathcal{M}[t]$ by

$$\partial_h p(t) = \frac{p(t) - p(t-h)}{h}, \quad \text{for } p \in \mathcal{M}[t].$$

It is clear that

$$\partial_h^n p(t) = \frac{1}{h^n} \sum_{j=0}^n (-1)^j \binom{n}{j} p(t-jh),$$

and, if $p = p_m t^m + \dots + p_0$, one has

$$\partial_h^n p(t) = \begin{cases} 0, & \text{if } n > m \\ m! p_m, & \text{if } n = m. \end{cases}$$

If $(t)_n$ is the Pochhammer polynomial of degree n , that is

$$(t)_0 = 1 \quad \text{and} \quad (t)_n = t(t+1) \dots (t+n-1) \quad \text{for } n \geq 1,$$

it is easy to see that $\partial_h(t/h)_n = n(t/h)_{n-1}$ and thus

$$\begin{aligned} \partial_h^r(t/h)_n &= r! \binom{n}{r} (t/h)_{n-r} \\ (2.4) \quad &= (n-r+1)_r (t/h)_{n-r} \quad \text{for } r \leq n. \end{aligned}$$

Given $r \in \mathcal{A}$ we extend to $\mathcal{M}[t]$ the adjoint action of r in $\mathcal{M}$ by

$$\text{ad}_r(p_0 + p_1 t + \dots + p_n t^n) = \text{ad}_r(p_0) + \text{ad}_r(p_1) t + \dots + \text{ad}_r(p_n) t^n.$$

Observe that ad_r commute with the discrete derivative ∂_h for all $h \in \mathbf{k}$. In addition, if $E, H \in \mathcal{A}$ are elements that satisfy $\text{ad}_H(E) = cE$, $c \in \mathbf{k}$, then it is straightforward to prove that

$$(2.5) \quad \text{ad}_E(p(t+H)) = \text{ad}_E(p)(t+H) - c \partial_c p(t+H) E$$

$$(2.6) \quad E^n p(t+H) = \sum_{j=0}^n \binom{n}{j} \text{ad}_E^{n-j}(p)(t-cj+H) E^j \quad \text{for all } n \in \mathbb{N}$$

$$(2.7) \quad E^n H^m = (H - nc)^m E^n \quad \text{for all } m, n \in \mathbb{N}.$$

2.3. The gaussian elimination. For $k \geq 0$ and $n \geq k$ we introduce the following equation,

$$\begin{aligned} \text{eq}_n^k : \sum_{i=0}^k \sum_{j=0}^{n-k} (-1)^i \binom{k}{i} \binom{n-k}{j} \text{ad}_E^{n-(j+i)} \partial_1^k \partial_{c-1}^i p(H + n - cj - i) E^{j+i} \\ = \sum_{i=0}^k (-1)^i \binom{k}{i} \text{ad}_E^{k-i} \partial_1^k \partial_{c-1}^i p(H - n + 2k - i) E^{n-(k-i)}. \end{aligned}$$

We now collect some properties of these equations.

- (1) The equation eq_n^0 is, $\sum_{j=0}^n \binom{n}{j} \text{ad}_E^{n-j} p(H + n - cj) E^j = p(H - n) E^n$. Hence, in view of the identity (2.6), the system $\{\text{eq}_n^0 : n \in \mathbb{N}\}$ is the original system (1.3).
- (2) On the other hand, the equation eq_n^n is

$$\begin{aligned} \sum_{i=0}^n (-1)^i \binom{n}{i} \text{ad}_E^{n-i} \partial_1^n \partial_{c-1}^i p(H + n - i) E^i \\ = \sum_{i=0}^n (-1)^i \binom{n}{i} \text{ad}_E^{n-i} \partial_1^n \partial_{c-1}^i p(H - n + 2n - i) E^i, \end{aligned}$$


which is trivial for all $n \in \mathbb{N}$.

- (3) The fundamental property of eq_n^k , for $k \geq 0$ and $n \geq k$, is that the polynomial p appears under a derivative of order greater than or equal to k . This implies that for all $n \geq k$ the equation eq_n^k involves only the coefficients p_j with $j \geq k$.

For $k \geq 0$, let Eq^k be the system

$$\text{Eq}^k = \{\text{eq}_1^0, \text{eq}_2^1, \dots, \text{eq}_k^{k-1}\} \cup \{\text{eq}_n^k : n > k\}.$$

As we have just observed, Eq^0 is the original system (1.3). The following picture may help to keep these definitions in mind.



Finally, let Eq^∞ be the system

$$\text{Eq}^\infty = \{\text{eq}_{n+1}^n : n \geq 0\}.$$

According to item (3) above, Eq^∞ is an upper triangular system, in the sense that if $p = \sum_{j \geq 0} p_j t^j$ then eq_{n+1}^n only involves the unknowns p_j with $j \geq n$. The equations of this system are the following,

$$\begin{aligned} \text{eq}_{n+1}^n : & \sum_{i=0}^n (-1)^i \binom{n}{i} \text{ad}_E^{n+1-i} \partial_1^n \partial_{c-1}^i p(H+n+1-i) E^i \\ & + \sum_{i=0}^n (-1)^i \binom{n}{i} \text{ad}_E^{n-i} \partial_1^n \partial_{c-1}^i p(H+n+1-c-i) E^{i+1} \\ & = \sum_{i=0}^n (-1)^i \binom{n}{i} \text{ad}_E^{n-i} \partial_1^n \partial_{c-1}^i p(H+n-1-i) E^{i+1}. \end{aligned}$$

Remark 2.1. If $\deg(p) = N$ then the only non zero equations of Eq^∞ are eq_{n+1}^n for $n = 0, \dots, N$. In this case, the last two non zero equations are,

$$\begin{aligned} \text{eq}_{N+1}^N : & \text{ad}_E^{N+1}(p_N) = 0, \\ \text{eq}_N^{N-1} : & \text{ad}_E^N(p_{N-1}) + N \text{ad}_E^N(p_N)(H + \frac{N+1}{2}) + N(N+1-Nc) \text{ad}_E^{N-1}(p_N)E = 0. \end{aligned}$$

Indeed, the equation eq_{N+1}^N is

$$\text{ad}_E^{N+1} \partial_1^N p(H+N+1) + \text{ad}_E^N \partial_1^N p(H+N+1-c)E = \text{ad}_E^N \partial_1^N p(H+N-1)E,$$

however, since $\partial_1^N p = N! p_N$ the equation eq_{N+1}^N becomes $\text{ad}_E^{N+1}(p_N) = 0$. For eq_N^{N-1} we use that $\partial_1^{N-1} p = (N-1)!(N p_N t - \binom{N}{2} p_N + p_{N-1})$, and proceed in a similar way.

The main theorem of this section is the following.

Theorem 2.2. *Assume that $[H, E] = cE$. For all $k \geq 0$, the first N equations of the system Eq^k are equivalent to the first N equations of the system Eq^0 . In particular this is true for the upper triangular system Eq^∞ .*

Proof. The theorem easily follows by induction once we prove that the equations eq_n^k satisfy the following recursion formula

$$(2.8) \quad \text{eq}_n^k = \text{eq}_n^{k-1} - 2 \text{eq}_{n-1}^{k-1} E + \text{eq}_{n-2}^{k-1} E^2 - \text{ad}_E(\text{eq}_{n-1}^{k-1}) + \text{ad}_E(\text{eq}_{n-2}^{k-1})E$$

for $n > k > 0$.

The proof of this recursion formula is very technical and will be done at the end of this section. We point out that this formula shows that, for $n > k$, the equation eq_n^k of Eq^k is equal to the equation eq_n^{k-1} of Eq^{k-1} plus a linear combination of the two previous equations of Eq^{k-1} . $\square$

Corollary 2.3. *Assume that $[H, E] = cE$. If $p \in \mathcal{M}[t]$ satisfies*

$$E^n p(H+n) = p(H-n) E^n \quad \text{for } n = 1, \dots, \deg(p),$$

then $E^n p(H+n) = p(H-n) E^n$ holds for all $n \geq 0$.

Proof. Let $N = \deg(p)$. The assumption on p says that it satisfies the first N equations of the system Eq^0 , then Theorem 2.2 implies that p satisfies the first N equations of Eq^∞ . On the other hand, we showed in Remark 2.1 that p automatically satisfies all the remaining equations of Eq^∞ . Therefore p satisfies all the equations of Eq^0 . $\square$

In the following corollary we describe the particular cases in which $c = 0$ and $c = 1$, in this cases the triangularized system became much simpler.

Corollary 2.4. *If $\text{ad}_H(E) = 0$ then the equations of the system Eq^∞ are*

$$\begin{aligned} \text{eq}_{n+1}^n : \sum_{i=0}^n \left[\binom{n}{i} \text{ad}_E^{n+1-i} \partial_1^{n+i} p(H+n+1)E^i + \binom{n}{i} \text{ad}_E^{n-i} \partial_1^{n+i} p(H+n+1)E^{i+1} \right] \\ = \sum_{i=0}^n \binom{n}{i} \text{ad}_E^{n-i} \partial_1^{n+i} p(H+n-1)E^{i+1}. \end{aligned}$$

If $\text{ad}_H(E) = E$ then the equations of the system Eq^∞ are

$$\text{eq}_{n+1}^n : \text{ad}_E^{n+1} \partial_1^n p(H+n+1) + \text{ad}_E^n \partial_1^{n+1} p(H+n)E = 0.$$

Proof. If E and H commute then $c = 0$, hence

$$\partial_{c-1} p(t) = \partial_{-1} p(t) = -\partial_1 p(t+1).$$

Therefore $\partial_{c-1}^i p(t) = (-1)^i \partial_1^i p(t+i)$ and the formula follows. On the other hand, if $\text{ad}_H(E) = E$ then $c = 1$, and hence $\partial_{c-1} = \partial_0 = 0$. Therefore all the terms of eq_{n+1}^n for $i > 0$ are zero and thus

$$\text{eq}_{n+1}^n : \text{ad}_E^{n+1} \partial_1^n p(H+n+1) + \text{ad}_E^n \partial_1^n p(H+n)E = \text{ad}_E^n \partial_1^n p(H+n-1)E,$$

which is equal to the equation stated in the corollary. $\square$

2.4. Proof of the Recursion Formula 2.8. We work first on the left hand sides (LHS) of the equations involved in the Recursion Formula 2.8 (see the definition of eq_n^k in Subsection 2.3). In fact, we need to prove that

$$\text{LHS}(\text{eq}_n^{k-1} - 2\text{eq}_{n-1}^{k-1} E + \text{eq}_{n-2}^{k-1} E^2 - \text{ad}_E(\text{eq}_{n-1}^{k-1}) + \text{ad}_E(\text{eq}_{n-2}^{k-1})E) = \text{LHS}(\text{eq}_n^k).$$

Since the sums in the definition of eq_n^k runs over all possible values of i and j for which the binomial numbers are not zero we omit them.

$$\begin{aligned}
& \text{LHS}(\text{eq}_n^{k-1}) : \\
& (-1)^i \binom{k-1}{i} \binom{n-k+1}{j} \text{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-cj-i) E^{j+i} \\
& \text{LHS}(-2 \text{eq}_{n-1}^{k-1} E) : \\
& -(-1)^i \binom{k-1}{i} \binom{n-k}{j} \text{ad}_E^{n-1-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-1-cj-i) E^{j+i+1} \\
& -(-1)^i \binom{k-1}{i} \binom{n-k}{j} \text{ad}_E^{n-1-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-1-cj-i) E^{j+i+1} \\
& \text{LHS}(+ \text{eq}_{n-2}^{k-1} E^2) : \\
& +(-1)^i \binom{k-1}{i} \binom{n-1-k}{j} \text{ad}_E^{n-2-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-2-cj-i) E^{j+i+2} \\
& \text{LHS}(-\text{ad}_E(\text{eq}_{n-1}^{k-1})) : \\
& -(-1)^i \binom{k-1}{i} \binom{n-k}{j} \text{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-1-cj-i) E^{j+i} \\
& -(-1)^i \binom{k-1}{i} \binom{n-k}{j} \text{ad}_E^{n-1-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-1-cj-c-i) E^{j+i+1} \\
& +(-1)^i \binom{k-1}{i} \binom{n-k}{j} \text{ad}_E^{n-1-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-1-cj-i) E^{j+i+1} \\
& \text{LHS}(+ \text{ad}_E(\text{eq}_{n-2}^{k-1} E)) : \\
& +(-1)^i \binom{k-1}{i} \binom{n-1-k}{j} \text{ad}_E^{n-1-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-2-cj-i) E^{j+i+1} \\
& +(-1)^i \binom{k-1}{i} \binom{n-1-k}{j} \text{ad}_E^{n-2-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-2-cj-c-i) E^{j+i+2} \\
& -(-1)^i \binom{k-1}{i} \binom{n-1-k}{j} \text{ad}_E^{n-2-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-2-cj-i) E^{j+i+2}
\end{aligned}$$

We now indicate how to simplify these ten lines.

Lines 3 and 7 cancel out and so do lines 4 and 10. Lines 6, 8 and 9, after a change of variables, become

$$\begin{aligned}
& -(-1)^i \binom{k-1}{i} \binom{n-k}{j-1} \text{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-1-cj-i) E^{j+i} \\
& +(-1)^i \binom{k-1}{i} \binom{n-1-k}{j-1} \text{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-2-cj+c-i) E^{j+i} \\
& +(-1)^i \binom{k-1}{i} \binom{n-1-k}{j-2} \text{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-2-cj+c-i) E^{j+i},
\end{aligned}$$

which is equal to

$$\begin{aligned}
& -(-1)^i \binom{k-1}{i} \binom{n-k}{j-1} \text{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-1-cj-i) E^{j+i} \\
& +(-1)^i \binom{k-1}{i} \binom{n-k}{j-1} \text{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^i b(H+n-2-cj+c-i) E^{j+i}
\end{aligned}$$

and finally equal to

$$(-1)^i \binom{k-1}{i} \binom{n-k}{j-1} \text{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^{i+1} b(H+n-1-c(j-1)-(i+1)) E^{j+i}.$$

Similarly, lines 1, 2 and 5 yield

$$\begin{aligned} & (-1)^i \binom{k-1}{i} \binom{n-k}{j} \operatorname{ad}_E^{n-(j+i)} \partial_1^k \partial_{c-1}^i b(H+n-cj-i) E^{j+i} \\ & - (-1)^i \binom{k-1}{i} \binom{n-k}{j-1} \operatorname{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^{i+1} b(H+n-c(j-1)-(i+1)) E^{j+i}. \end{aligned}$$

Thus we have,

$$\begin{aligned} & (-1)^i \binom{k-1}{i} \binom{n-k}{j-1} \operatorname{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^{i+1} b(H+n-1-c(j-1)-(i+1)) E^{j+i} \\ & + (-1)^i \binom{k-1}{i} \binom{n-k}{j} \operatorname{ad}_E^{n-(j+i)} \partial_1^k \partial_{c-1}^i b(H+n-cj-i) E^{j+i} \\ & - (-1)^i \binom{k-1}{i} \binom{n-k}{j-1} \operatorname{ad}_E^{n-(j+i)} \partial_1^{k-1} \partial_{c-1}^{i+1} b(H+n-c(j-1)-(i+1)) E^{j+i} \\ = & - (-1)^i \binom{k-1}{i} \binom{n-k}{j-1} \operatorname{ad}_E^{n-(j+i)} \partial_1^k \partial_{c-1}^{i+1} b(H+n-c(j-1)-(i+1)) E^{j+i} \\ & + (-1)^i \binom{k-1}{i} \binom{n-k}{j} \operatorname{ad}_E^{n-(j+i)} \partial_1^k \partial_{c-1}^i b(H+n-cj-i) E^{j+i} \\ = & (-1)^i \binom{k-1}{i-1} \binom{n-k}{j} \operatorname{ad}_E^{n-(j+i)} \partial_1^k \partial_{c-1}^i b(H+n-cj-i) E^{j+i} \\ & + (-1)^i \binom{k-1}{i} \binom{n-k}{j} \operatorname{ad}_E^{n-(j+i)} \partial_1^k \partial_{c-1}^i b(H+n-cj-i) E^{j+i} \\ = & (-1)^i \binom{k}{i} \binom{n-k}{j} \operatorname{ad}_E^{n-(j+i)} \partial_1^k \partial_{c-1}^i b(H+n-cj-i) E^{j+i} \\ = & \text{LHS}(\text{eq}_n^k). \end{aligned}$$

We now work on the right hand sides (RHS) of the equations involved in the Recursion Formula 2.8.

$$\text{RHS}(\text{eq}_n^{k-1}) :$$

$$(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+2(k-1)-i) E^{n-(k-1-i)}$$

$$\text{RHS}(-2 \text{eq}_{n-1}^{k-1} E) :$$

$$-(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+1+2(k-1)-i) E^{n-(k-1-i)}$$

$$-(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+1+2(k-1)-i) E^{n-(k-1-i)}$$

$$\text{RHS}(+ \text{eq}_{n-2}^{k-1} E^2) :$$

$$+(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+2+2(k-1)-i) E^{n-(k-1-i)}$$

$$\text{RHS}(- \operatorname{ad}_E(\text{eq}_{n-1}^{k-1})) :$$

$$-(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+1+2(k-1)-i) E^{n-(k-i)}$$

$$-(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+1+2(k-1)-c-i) E^{n-(k-1-i)}$$

$$+(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+1+2(k-1)-i) E^{n-(k-1-i)}$$

$$\text{RHS}(+ \operatorname{ad}_E(\text{eq}_{n-2}^{k-1}) E) :$$

$$+(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+2+2(k-1)-i) E^{n-(k-i)}$$

$$+(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+2+2(k-1)-c-i) E^{n-(k-1-i)}$$

$$-(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^{k-1} \partial_{c-1}^i b(H-n+2+2(k-1)-i) E^{n-(k-1-i)}$$

Now the sum of lines 1 and 2 is equal to

$$-(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^k \partial_{c-1}^i b(H - n - 1 + 2k - i) E^{n-(k-1-i)},$$

similarly the sum of lines 5 and 8 is equal to

$$(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-i} \partial_1^k \partial_{c-1}^i b(H - n + 2k - i) E^{n-(k-i)},$$

and the sum of lines 6 and 9 is equal to

$$(-1)^i \binom{k}{i} \operatorname{ad}_E^{k-1-i} \partial_1^k \partial_{c-1}^i b(H - n + 2k - c - i) E^{n-(k-1-i)}.$$

Finally, the sum of the last three lines is equal to

$$\begin{aligned} & -(-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-1-i} \partial_1^k \partial_{c-1}^{i+1} b(H - n + 2k - (i+1)) E^{n-(k-1-i)} \\ & + (-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-i} \partial_1^k \partial_{c-1}^i b(H - n + 2k - i) E^{n-(k-i)} \\ & = (-1)^i \binom{k-1}{i-1} \operatorname{ad}_E^{k-i} \partial_1^k \partial_{c-1}^i b(H - n + 2k - i) E^{n-(k-i)} \\ & \quad + (-1)^i \binom{k-1}{i} \operatorname{ad}_E^{k-i} \partial_1^k \partial_{c-1}^i b(H - n + 2k - i) E^{n-(k-i)} \\ & = (-1)^i \binom{k}{i} \operatorname{ad}_E^{k-i} \partial_1^k \partial_{c-1}^i b(H - n + 2k - i) E^{n-(k-i)} \\ & = \operatorname{RHS}(\operatorname{eq}_n^k). \end{aligned}$$

This finishes the proof of the Recursion Formula 2.8.

The following two sections are devoted to express the system (1.3) in a matrix form and to give its LU-decomposition in a general context. In particular we will not assume that $[H, E] = cE$.

3. INFINITE MATRICES

AND THE MATRIX REPRESENTATION OF THE SYSTEM (1.3)

In this section we recall some preliminaries on infinite matrices, introduce the concept of *periodic Gaussian elimination process* and present the matrix representation of the system (1.3).

3.1. Infinite matrices. Given an associative k -algebra $\mathcal{A}$, let $\mathcal{M}_\infty(\mathcal{A})$ denote the set of all infinite matrices

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdot \\ a_{21} & a_{22} & a_{23} & \cdot \\ a_{31} & a_{32} & a_{33} & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

with $a_{ij} \in \mathcal{A}$. The n -minor of a matrix $A \in \mathcal{M}_\infty(\mathcal{A})$ is the matrix corresponding to the upper-left corner of A of size $n \times n$. We say that a sequence of matrices B_k converges to B if for every i, j there exists k_0 such that $(B_k)_{ij} = B_{ij}$ for all $k \geq k_0$. A matrix $A \in \mathcal{M}_\infty(\mathcal{A})$ is said *row-finite* (resp. *column-finite*) if every row (resp. column) of A contains only a finite number of non zero elements. Lower triangular and upper triangular matrices are, respectively, examples of row-finite and column-finite matrices.

It is clear that $\mathcal{M}_\infty(\mathcal{A})$ is not a ring since the multiplication of two matrices does not always exist. Nevertheless, if $A, B \in \mathcal{M}_\infty(\mathcal{A})$ and either A is

row-finite or B is column-finite then AB do exist. It is not difficult to prove the following proposition.

Proposition 3.1.

- (a) *A lower or upper triangular matrix $A \in \mathcal{M}_\infty(\mathcal{A})$ is invertible if and only if all n -minors of A are invertible.*
- (b) *An LU-factorization of a matrix $A \in \mathcal{M}_\infty(\mathcal{A})$ exists if and only if it exists for all n -minors of A . Moreover, in this case the LU-factorization of A is unique.*

Definition 3.2. Given a matrix $A \in \mathcal{M}_\infty(\mathcal{A})$ the *shifted matrix* $s(A)$ of A is the diagonal blocked matrix formed by the identity matrix of size 1×1 and A , that is

$$s(A) = \begin{pmatrix} 1 & \\ & A \end{pmatrix}.$$

To apply an elementary row operation to a matrix A is equivalent to multiply A on the left by a lower triangular matrix. For some special matrices $A \in \mathcal{M}_\infty(\mathcal{A})$ it is possible to perform the elimination process of Gauss by using a periodic sequence of elementary row operations, in the sense that there exist a lower triangular matrix $T_0 \in \mathcal{M}_\infty(\mathcal{A})$, that would constitute the fundamental part of the periodic sequence, such that the whole periodic sequence of elementary row operations is of the form,

$$T_0; \text{ next } s(T_0); \text{ next } s^2(T_0); \text{ next } s^3(T_0); \dots$$

In other words, this means that the sequence of matrices

$$A, \quad T_0 A, \quad s(T_0) T_0 A, \quad s^2(T_0) s(T_0) T_0 A, \quad s^3(T_0) s^2(T_0) s(T_0) T_0 A, \quad \dots$$

converges to an upper triangular matrix. This suggest the following definitions.

Definition 3.3. Given a lower triangular matrix $T_0 \in \mathcal{M}_\infty(\mathcal{A})$ the *left iterated matrix* T_0^L corresponding to T_0 is the infinite (from right to left) product

$$T_0^L = \dots s^3(T_0) s^2(T_0) s(T_0) T_0.$$

It is clear that this product converges to a lower triangular matrix. Similarly, the *right iterated matrix* T_0^R corresponding to T_0 is the lower triangular matrix given by the infinite (from left to right) product

$$T_0^R = T_0 s(T_0) s^2(T_0) s^3(T_0) \dots$$

We say that a matrix $A \in \mathcal{M}_\infty(\mathcal{A})$ admits a *periodic Gaussian elimination process* if there exist a lower triangular matrix $T_0 \in \mathcal{M}_\infty(\mathcal{A})$ such that $T_0^L A$ is upper triangular. In this case T_0 is called a *fundamental sequence* of the Gaussian elimination process of A .

It is not difficult to prove that T_0^L and T_0^R are respectively characterized by the identities

$$T_0^L = s(T_0^L) T_0 \quad \text{and} \quad T_0^R = T_0 s(T_0^R),$$

and it is clear that if T_0 is invertible then T_0^L is also invertible and

$$(T_0^L)^{-1} = (T_0^{-1})^R.$$

Now the following proposition is clear.

Proposition 3.4. *An infinite matrix A admits a periodic gaussian elimination process if and only if A admits an LU -decomposition. In fact, if $A = LU$ then the fundamental period is $T_0 = s(L)L^{-1}$ and, conversely, if T_0 is the fundamental period of A then*

$$L^{-1} = (T_0)^L \text{ and } L = (T_0^{-1})^R.$$

3.2. Relevant infinite matrices. In this subsection we introduce some infinite matrices that will be used frequently in what follows and we also collect some nice properties that these matrices enjoy.

- . The Vandermonde matrix: $V_{ij} = i^{j-1}$.
- . The diagonal matrix formed by the powers of $q \in \mathcal{A}$: $(D_q)_{ij} = \delta_{ij} q^i$.
- . The diagonal matrix formed by the factorial numbers: $F_{ij} = \delta_{ij} (i-1)!$.
- . The lower triangular matrix formed by the Pascal numbers: $P_{ij} = \binom{i-1}{j-1}$.
- . The lower triangular matrix formed by the Stirling numbers of the second kind: $S_{ij} = S(i, j)$ for $i \geq j$, where $S(i, j) = \frac{1}{j!} \sum_{k=0}^j (-1)^k \binom{j}{k} (j-k)^i$.

These matrices are,

$$V = \begin{pmatrix} 1 & 1 & 1 & 1 & \cdot \\ 1 & 2 & 4 & 8 & \cdot \\ 1 & 3 & 9 & 27 & \cdot \\ 1 & 4 & 16 & 64 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}, \quad D_q = \begin{pmatrix} q^0 & & & & \\ & q^2 & & & \\ & & q^3 & & \\ & & & q^4 & \\ & & & & \cdot \end{pmatrix}, \quad F = \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 2 & & \\ & & & 6 & \\ & & & & \cdot \end{pmatrix},$$

$$P = \begin{pmatrix} 1 & & & & \cdot \\ 1 & 1 & & & \cdot \\ 1 & 2 & 1 & & \cdot \\ 1 & 3 & 3 & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}, \quad S = \begin{pmatrix} 1 & & & & \cdot \\ 1 & 1 & & & \cdot \\ 1 & 3 & 1 & & \cdot \\ 1 & 7 & 6 & 1 & \cdot \\ 1 & 15 & 25 & 10 & 1 \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

Next we collect some of the properties of these matrices:

- (1) The Vandermonde matrix V has an LDU -factorization. It is given by $V = PFS^t$.
- (2) $P^{-1} = D_{-1}PD_{-1}$.
- (3) The classical recurrence relations

$$\binom{i}{j} = \binom{i-1}{j-1} + \binom{i-1}{j} \quad \text{and} \quad S(i, j) = S(i-1, j-1) + jS(i-1, j)$$

that define, respectively, the Pascal and Stirling numbers correspond to the matrix identities $P = s(P)T_{0,P}$ and $S = s(S)T_{0,S}$ where

$$T_{0,P} = \begin{pmatrix} 1 & 1 & & \cdot \\ 1 & 1 & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix} \quad \text{and} \quad T_{0,S} = \begin{pmatrix} 1 & 1 & & \cdot \\ 1 & 2 & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

Thus P and S are the left iterated matrices $P = T_{0,P}^L$ and $S = T_{0,S}^L$.

- (4) P and S are also right iterated matrices. Since $P = \begin{pmatrix} 1 & 1 & & \cdot \\ 1 & 1 & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix} s(P)$ and

$$S = P s(S) \text{ we have } P = \begin{pmatrix} 1 & 1 & & \cdot \\ 1 & 1 & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix}^R \text{ and } S = P^R.$$

where

$$P_x = \left(\binom{i-1}{j-1} x^{i-j} \right) = \begin{pmatrix} 1 & & & & \\ x & 1 & & & \\ x^2 & 2x & 1 & & \\ x^3 & 3x^2 & 3x & 1 & \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Hence $M = M_0 P_{R_H}^t$ where

$$M_0 = \begin{pmatrix} L_E - R_E & L_E + R_E & L_E - R_E & L_E + R_E & \cdot \\ L_E^2 - R_E^2 & 2L_E^2 + 2R_E^2 & 4L_E^2 - 4R_E^2 & 8L_E^2 + 8R_E^2 & \cdot \\ L_E^3 - R_E^3 & 3L_E^3 + 3R_E^3 & 9L_E^3 - 9R_E^3 & 27L_E^3 + 27R_E^3 & \cdot \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

It is also convenient to consider the Pochhammer basis $\{(t)_j\}$ of $k[t]$, instead of the basis $\{t^j\}$, to express the system (1.3). If we do this, an analysis similar to the previous one shows that $p = a_0(t)_0 + a_1(t)_1 + a_2(t)_2 + \dots \in \mathcal{M}[t]$ satisfies the system (1.3) if and only if $M' \begin{pmatrix} a_0 \\ a_1 \\ a_2 \\ \vdots \end{pmatrix} = 0$, where

$$M' = (M'_1 - M'_2)F(P'_{R_H})^t,$$

and the matrices M'_1 , M'_2 and P'_x are as follows,

$$P'_x = \left(\binom{i-1}{j-1} (x)_{i-j} \right) = \begin{pmatrix} 1 & & & & \\ (x)_1 & 1 & & & \\ (x)_2 & 2(x)_1 & 1 & & \\ (x)_3 & 3(x)_2 & 3(x)_1 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

$$M'_1 = D_{L_E} \begin{pmatrix} 1 & 1 & 1 & 1 & \cdot \\ 1 & 2 & 3 & 4 & \cdot \\ 1 & 3 & 6 & 10 & \cdot \\ 1 & 4 & 10 & 20 & \cdot \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad \text{and} \quad M'_2 = D_{R_E} \begin{pmatrix} 1 & -1 & & & \\ 1 & -2 & 1 & & \\ 1 & -3 & 3 & -1 & \\ 1 & -4 & 6 & -4 & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Remark 3.5. According to item (6) of §3.2, the matrices M and M' are related by

$$M = M' D_{-1} s(S)^t D_{-1}.$$

Thus, the matrix M' could be seen as the result of factoring out from M the Stirling numbers that the Vandermonde matrix (appearing in M) contains.

4. THE LU-DECOMPOSITION OF THE SYSTEM (1.3)

4.1. The L and U factors. We have just seen that M is the matrix corresponding to system (1.3) in the canonical basis and M' is the corresponding matrix in the Pochhammer basis. We also have

$$M = M_0 P_{R_H}^t, \quad M' = (M'_1 - M'_2)F(P'_{R_H})^t, \quad M = M' D_{-1} s(S)^t D_{-1}.$$

Therefore, in order to find the LU-decomposition of the system (1.3) it is sufficient to find the LU-decomposition of either M_0 or $M'_1 - M'_2$. Moreover, the entries of M_0 and $M'_1 - M'_2$ are homogeneous polynomials in the variables L_E and R_E . Since an homogeneous polynomial $q(x_1, x_2)$ in two variables x_1 and x_2 is completely determined by the polynomial $\tilde{q}(x) = q(x, 1)$, we shall simplify the notation by substituting the variables

$$(4.1) \quad L_E \text{ by } x \quad \text{and} \quad R_E \text{ by } 1.$$

Under this transformations, the matrices M_0 , M'_1 and M'_2 become

$$\tilde{M}_0 = \begin{pmatrix} x-1 & x+1 & x-1 & x+1 & \cdot \\ x^2-1 & 2x^2+2 & 4x^2-4 & 8x^2+8 & \cdot \\ x^3-1 & 3x^3+3 & 9x^3-9 & 27x^3+27 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix},$$

$$\tilde{M}'_1 = D_x \begin{pmatrix} 1 & 1 & 1 & 1 & \cdot \\ 1 & 2 & 3 & 4 & \cdot \\ 1 & 3 & 6 & 10 & \cdot \\ 1 & 4 & 10 & 20 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix} \quad \text{and} \quad \tilde{M}'_2 = \begin{pmatrix} 1 & -1 & & & \cdot \\ 1 & -2 & 1 & & \cdot \\ 1 & -3 & 3 & -1 & \cdot \\ 1 & -4 & 6 & -4 & 1 \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

We point out that the LU-decompositions of the matrices $\tilde{M}'_1$ and $\tilde{M}'_2$, separately, are

$$\tilde{M}'_1 = D_x P P^t \quad \text{and} \quad \tilde{M}'_2 = P \begin{pmatrix} -1 & 1 & & & \cdot \\ & 1 & -1 & & \cdot \\ & & -1 & 1 & \cdot \\ & & & 1 & -1 \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

The LU-decomposition of $\tilde{M}'_1 - \tilde{M}'_2$ is more subtle and it is obtained from Theorem 2.2 as follows:

- (1) $(\tilde{M}'_1 - \tilde{M}'_2)F$ is the matrix corresponding to the system (1.3) for $H = 0$, when it is expressed in terms of the basis $\{(t)_j\}$.
- (2) In other words, $(\tilde{M}'_1 - \tilde{M}'_2)F$ is the matrix corresponding to the system Eq^0 for $H = 0$, when it is expressed in terms of the basis $\{(t)_j\}$.
- (3) The definition of the system Eq^k and the Recursion Formula 2.8 imply that the system Eq^k is obtained by keeping the first k equations of the system Eq^{k-1} and by incorporating the equations

$$\text{eq}_i^k = \text{eq}_i^{k-1} - 2R_E(\text{eq}_{i-1}^{k-1}) + R_E^2(\text{eq}_{i-2}^{k-1}) - \text{ad}_E(\text{eq}_{i-1}^{k-1}) + R_E \text{ad}_E(\text{eq}_{i-2}^{k-1})$$

for $i > k > 0$. Applying the change of variables (4.1) these equations are transformed as follows,

$$\text{eq}_i^k = \text{eq}_i^{k-1} - (x+1)\text{eq}_{i-1}^{k-1} + x\text{eq}_{i-2}^{k-1}.$$

This means that the lower triangular matrix that transforms the system Eq^{k-1} into the system Eq^k is $s^{k-1}(T_0)$ with

$$(4.2) \quad T_0 = \begin{pmatrix} 1 & & & & \cdot \\ -x-1 & 1 & & & \cdot \\ x & -x-1 & 1 & & \cdot \\ & x & -x-1 & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

- (4) Theorem 2.2 implies that the left iterated matrix T_0^L associated to T_0 transforms the system Eq^0 into the system Eq^∞ .
- (5) Let

$$\tilde{M}'_1 - \tilde{M}'_2 = \tilde{L}'\tilde{U}'$$

be the LU-decomposition of $\tilde{M}'_1 - \tilde{M}'_2$. According to the previous item, we have that $\tilde{L}' = (T_0^L)^{-1} = (T_0^{-1})^R$ and $\tilde{U}'$ is of the form $\tilde{U}'_1 - \tilde{U}'_2$ where $\tilde{U}'_1 F$ (respectively, $\tilde{U}'_2 F$) is the matrix corresponding to the left (respectively, right) hand side of the system Eq^∞ for $H = 0$ (and thus $c = 0$), when the system is expressed in terms of the basis $\{(t)_j\}$.

(6) According to Corollary 2.4, when $H = 0$, the i th equation of Eq^∞ is

$$\begin{aligned} \text{eq}_i^{i-1} : \sum_{r=0}^{i-1} \left[\binom{i-1}{r} \text{ad}_E^{i-r} \partial_1^{i+r-1} p(i) E^r + \binom{i-1}{r} \text{ad}_E^{i-r-1} \partial_1^{i+r-1} p(i) E^{r+1} \right] \\ = \sum_{r=0}^{i-1} \binom{i-1}{r} \text{ad}_E^{i-r-1} \partial_1^{i+r-1} p(i-2) E^{r+1}. \end{aligned}$$

Now, replacing ad_E by $(x-1)$ and E by 1 we obtain,

$$\begin{aligned} \text{eq}_i^{i-1} : \sum_{r=0}^{i-1} \binom{i-1}{r} \partial_1^{i+r-1} p(i) x (x-1)^{i-r-1} \\ = \sum_{r=0}^{i-1} \binom{i-1}{r} \partial_1^{i+r-1} p(i-2) (x-1)^{i-r-1}. \end{aligned}$$

(7) If $p = a_0(t)_0 + a_1(t)_1 + a_2(t)_2 + \dots$ then, applying (2.4), we obtain

$$\begin{aligned} \text{eq}_i^{i-1} : \sum_{j \geq 0} \sum_{r=0}^{j-i+1} \binom{i-1}{r} (j-i-r+2)_{i+r-1} (i)_{j-i-r+1} x (x-1)^{i-r-1} a_j \\ = \sum_{j \geq 0} \sum_{r=0}^{j-i+1} \binom{i-1}{r} (j-i-r+2)_{i+r-1} (i-2)_{j-i-r+1} (x-1)^{i-r-1} a_j. \end{aligned}$$

(8) Now, the (i, j) entry of the matrix $\tilde{U}'_1 F$ (respectively, $\tilde{U}'_2 F$) is the coefficient multiplying a_{j-1} in the left (respectively, right) hand side of the i th equation the system Eq^∞ given in (7), that is

$$\begin{aligned} (\tilde{U}'_1)_{ij} &= \frac{1}{(j-1)!} \sum_{r=0}^{j-i} \binom{i-1}{r} (j-i-r+1)_{i+r-1} (i)_{j-i-r} (x-1)^{i-r-1} x, \\ (\tilde{U}'_2)_{ij} &= \frac{1}{(j-1)!} \sum_{r=0}^{j-i} \binom{i-1}{r} (j-i-r+1)_{i+r-1} (i-2)_{j-i-r} (x-1)^{i-r-1}. \end{aligned}$$

We shall now complete the description of the LU-decomposition of the system (1.3) by showing that $\tilde{L}'$, $(\tilde{L}')^{-1}$, $\tilde{U}'_1$ and $\tilde{U}'_2$ are matrices whose entries are given by some particular Jacobi polynomials. The basic facts about Jacobi polynomials that we shall need are collected in §5.

4.2. The entries of L and L^{-1} are ultraspherical Jacobi polynomials.

Recall that according to item(5) of §4.1 we know that $\tilde{L}' = (T_0^{-1})^R$ and $(\tilde{L}')^{-1} = T_0^L$. It is easy to see that the inverse of

$$T_0 = \begin{pmatrix} 1 & & & \\ -x-1 & 1 & & \\ x & -x-1 & 1 & \\ & x & -x-1 & 1 \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

is the lower triangular matrix with entries $(T_0^{-1})_{ij} = \frac{x^{i-j+1}-1}{x-1}$ if $i \geq j$, that is

$$T_0^{-1} = \begin{pmatrix} 1 & & & \\ 1+x & 1 & & \\ 1+x+x^2 & 1+x & 1 & \\ 1+x+x^2+x^3 & 1+x+x^2 & 1+x & 1 \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix}.$$

Theorem 4.1. *The entries of $\tilde{L}'$ and its inverse $(\tilde{L}')^{-1}$ are given by the following ultraspherical Jacobi polynomials.*

$$(\tilde{L}')_{ij} = (-1)^{i-j} p_{i-j}^{-i,-i}(x),$$

$$\left((\tilde{L}')^{-1}\right)_{ij} = (-1)^{i-j} \frac{j}{i} p_{i-j}^{j,j}(x).$$

Proof. Let K' and K'' be the matrices defined by the right hand sides of the above equalities, that is

$$K'_{ij} = (-1)^{i-j} p_{i-j}^{-i,-i}(x) \quad \text{and} \quad K''_{ij} = (-1)^{i-j} \frac{j}{i} p_{i-j}^{j,j}(x).$$

By the definition of left and right iterated matrices, it is sufficient to prove that

$$K' = T_0^{-1} s(K') \quad \text{and} \quad K'' = s(K'') T_0.$$

The first identity holds if and only if

$$(-1)^{i-j} p_{i-j}^{-i,-i}(x) = \sum_{k=j}^i \frac{x^{i-k+1} - 1}{x - 1} (-1)^{k-j} p_{k-j}^{-k+1,-k+1}(x),$$

or equivalently, if the following equality holds

$$p_{i-j}^{-i,-i}(x) = \sum_{k=0}^{i-j} \frac{x^{i-j-k+1} - 1}{x - 1} (-1)^{k+j-i} p_k^{-k-j+1,-k-j+1}(x),$$

which in turn holds if and only if

$$P_{i-j}^{-i,-i}\left(\frac{x+1}{x-1}\right) = \sum_{k=0}^{i-j} \left(1 - x^{i-j-k+1}\right) (1-x)^{k+j-i-1} P_k^{-k-j+1,-k-j+1}\left(\frac{x+1}{x-1}\right).$$

Now, this last identity is equivalent to

$$P_{i-j}^{-i,-i}(x) = \sum_{k=0}^{i-j+1} \left(\left(\frac{1-x}{2}\right)^{i-j-k+1} - \left(\frac{-1-x}{2}\right)^{i-j-k+1} \right) P_k^{-k-j+1,-k-j+1}(x),$$

and this last equality is true, since it is the difference of the two identities stated in Lemma 5.1 for $n = i - j + 1$ and $\alpha = i$. This completes the proof of the identity $K' = T_0^{-1} s(K')$.

The second identity holds if and only if

$$(-1)^{i-j} \frac{j}{i} p_{i-j}^{j,j}(x) = \sum_{k=j}^i (-1)^{i-k} \frac{k-1}{i-1} p_{i-k}^{k-1,k-1}(x) (T_0)_{kj}$$

for $i \geq 2$, or equivalently if

$$\frac{j}{i} p_{i-j}^{j,j}(x) = \frac{j-1}{i-1} p_{i-j}^{j-1,j-1}(x) + \frac{j(x+1)}{i-1} p_{i-j-1}^{j,j}(x) + \frac{(j+1)x}{i-1} p_{i-j-2}^{j+1,j+1}(x),$$

which in turn holds if and only if

$$\begin{aligned} & \frac{j}{i} P_{i-j}^{j,j}\left(\frac{x+1}{x-1}\right) \\ &= \frac{j-1}{i-1} P_{i-j}^{j-1,j-1}\left(\frac{x+1}{x-1}\right) + \frac{j}{i-1} \frac{x+1}{x-1} P_{i-j-1}^{j,j}\left(\frac{x+1}{x-1}\right) + \frac{j+1}{i-1} \frac{x}{(x-1)^2} P_{i-j-2}^{j+1,j+1}\left(\frac{x+1}{x-1}\right), \end{aligned}$$

or equivalently, if the following equality holds

$$4(i-1)j P_{i-j}^{j,j}(x) = 4i(j-1) P_{i-j}^{j-1,j-1}(x) + 4ijx P_{i-j-1}^{j,j}(x) + i(j+1)(x^2-1) P_{i-j-2}^{j+1,j+1}(x).$$

Now, this last identity is true since it is $(1+j)$ times equation (5.5) with $n = i-j-1$ and $\alpha = j+1$, plus $2(j-1)$ times equation (5.2) with $n = i-j$ and $\alpha = j$. This completes the proof of the identity $K'' = s(K'')T_0$. $\square$

Remark 4.2. It is known that the ultraspherical Jacobi polynomials constitute a 2-parameter family of polynomials that are orthogonal with respect to a continuous measure. The above theorem provides the following “discrete orthogonality” relationship that involves once many of them,

$$\begin{pmatrix} \frac{1}{1}p_0^{1,1} & & & \\ \frac{1}{2}p_1^{1,1} & \frac{2}{2}p_0^{2,2} & & \\ \frac{1}{3}p_2^{1,1} & \frac{2}{3}p_1^{2,2} & \frac{3}{3}p_0^{3,3} & \\ \frac{1}{4}p_3^{1,1} & \frac{2}{4}p_2^{2,2} & \frac{3}{4}p_1^{3,3} & \frac{1}{1}p_0^{4,4} \end{pmatrix} \begin{pmatrix} p_0^{-1,-1} & & & \\ p_1^{-2,-2} & p_0^{-2,-2} & & \\ p_2^{-3,-3} & p_1^{-3,-3} & p_0^{-3,-3} & \\ p_3^{-4,-4} & p_2^{-4,-4} & p_1^{-4,-4} & p_0^{-4,-4} \end{pmatrix} = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix}.$$

4.3. The entries of $\tilde{U}'_1$ and $\tilde{U}'_2$ are Jacobi polynomials.

Theorem 4.3. *The entries of $\tilde{U}'_1$ and $\tilde{U}'_2$ are given by the following Jacobi polynomials.*

$$\begin{aligned} (\tilde{U}'_1)_{ij} &= (-1)^{j-i} (x-1)^{2i-j-1} x p_{j-i}^{-j,0}(x), \\ (\tilde{U}'_2)_{ij} &= (-1)^{j-i} (x-1)^{2i-j-1} p_{j-i}^{-j+2,-2}(x). \end{aligned}$$

Proof. The theorem will be proved if we show that

$$\tilde{U}'_1 \left(\frac{x+1}{x-1} \right)_{ij} = (-1)^{j-i} \left(\frac{x-1}{2} \right)^{1-i} \frac{x+1}{x-1} P_{j-i}^{-j,0}(x),$$

and

$$\tilde{U}'_2 \left(\frac{x+1}{x-1} \right)_{ij} = (-1)^{j-i} \left(\frac{x-1}{2} \right)^{1-i} P_{j-i}^{-j+2,-2}(x).$$

In fact, according to item (8)

$$\begin{aligned} \tilde{U}'_1 \left(\frac{x+1}{x-1} \right)_{ij} &= \frac{x+1}{2(j-1)!} \sum_{r=0}^{j-i} \binom{i-1}{r} (j-i-r+1)_{i+r-1} (i)_{j-i-r} \left(\frac{x-1}{2} \right)^{r-i} \\ &= \frac{x+1}{2} \left(\frac{x-1}{2} \right)^{-i} \frac{1}{(j-i)!} \sum_{r=0}^{j-i} \binom{j-i}{r} (i-r)_r (i)_{j-i-r} \left(\frac{x-1}{2} \right)^r \\ &= \frac{x+1}{2} \left(\frac{x-1}{2} \right)^{-i} \frac{(-1)^{j-i}}{(j-i)!} \sum_{r=0}^{j-i} \binom{j-i}{r} (1-i)_r (1-j+r)_{j-i-r} \left(\frac{x-1}{2} \right)^r \\ &= (-1)^{j-i} \left(\frac{x-1}{2} \right)^{1-i} \frac{x+1}{x-1} P_{j-i}^{-j,0}(x). \end{aligned}$$

Analogously,

$$\begin{aligned}
\tilde{U}'_2 \left(\frac{x+1}{x-1} \right)_{ij} &= \frac{1}{(j-1)!} \sum_{r=0}^{j-i} \binom{i-1}{r} (j-i-r+1)_{i+r-1} (i-2)_{j-i-r} \left(\frac{x-1}{2} \right)^{r-i+1} \\
&= \left(\frac{x-1}{2} \right)^{1-i} \frac{1}{(j-i)!} \sum_{r=0}^{j-i} \binom{j-i}{r} (i-r)_r (i-2)_{j-i-r} \left(\frac{x-1}{2} \right)^r \\
&= \left(\frac{x-1}{2} \right)^{1-i} \frac{(-1)^{j-i}}{(j-i)!} \sum_{r=0}^{j-i} \binom{j-i}{r} (1-i)_r (3-j+r)_{j-i-r} \left(\frac{x-1}{2} \right)^r \\
&= (-1)^{j-i} \left(\frac{x-1}{2} \right)^{1-i} P_{j-i}^{-j+2, -2}(x).
\end{aligned}$$

This completes the proof of the theorem. $\square$

We have therefore proved the following theorem

Theorem 4.4. *The LU-decomposition of*

$$\tilde{M}'_1 - \tilde{M}'_2 = \begin{pmatrix} x-1 & x+1 & x & x & \cdot \\ x^2-1 & 2x^2+2 & 3x^2-1 & 4x^2 & \cdot \\ x^3-1 & 3x^3+3 & 6x^3-3 & 10x^3+1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

is $\tilde{M}'_1 - \tilde{M}'_2 = \tilde{L}'\tilde{U}'$ where

$$\begin{aligned}
(\tilde{L}')_{ij} &= (-1)^{i-j} p_{i-j}^{-i, -i}(x), & i \geq j; \\
(\tilde{U}')_{ij} &= (-1)^{j-i} (x-1)^{2i-j-1} \left(x p_{j-i}^{-j, 0}(x) - p_{j-i}^{-j+2, -2}(x) \right), & i \leq j.
\end{aligned}$$

Corollary 4.5. *The LU-decomposition of*

$$\tilde{M}_0 = \begin{pmatrix} x-1 & x+1 & x-1 & x+1 & \cdot \\ x^2-1 & 2x^2+2 & 4x^2-4 & 8x^2+8 & \cdot \\ x^3-1 & 3x^3+3 & 9x^3-9 & 27x^3+27 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

is

$$\tilde{M}_0 = \tilde{L}_0 \tilde{U}_0 \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 2 & & \\ & & & 6 & \\ & & & & 24 \end{pmatrix} \begin{pmatrix} 1 & -1 & 1 & -1 & 1 & \cdot \\ & 1 & -3 & 7 & -15 & \cdot \\ & & 1 & -6 & 25 & \cdot \\ & & & 1 & -10 & \cdot \\ & & & & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

where

$$\begin{aligned}
(\tilde{L}_0)_{ij} &= (-1)^{i-j} p_{i-j}^{-i, -i}(x), & i \geq j; \\
(\tilde{U}_0)_{ij} &= (-1)^{i-j} \frac{j}{i} (x-1)^{2i-j} p_{j-i}^{-j, -1}(x), & i \leq j;
\end{aligned}$$

and the explicit numerical matrices are respectively formed by the factorial and (up to the minus signs) Stirling numbers of the second kind.

Proof. We know that

$$\begin{aligned}
\tilde{M}_0 &= (\tilde{M}'_1 - \tilde{M}'_2) F D_{-1} s(S)^t D_{-1} \\
&= \tilde{L}' \tilde{U}' F D_{-1} s(S)^t D_{-1}.
\end{aligned}$$

This implies that $\tilde{L}'$ is the L-part of $\tilde{M}_0$ and thus we obtain the expression for $(\tilde{L}_0)_{ij}$. Now we must prove that

$$\tilde{U}_0 F D_{-1} S^t D_{-1} = \tilde{U}' F D_{-1} s(S)^t D_{-1}$$

which is easily seen to be equivalent to prove that

$$\tilde{U}_0 \begin{pmatrix} 1 & -1 & \cdot \\ & 1 & -1 \\ & & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix} = \tilde{U}'$$

that is

$$(x-1) \left(\frac{j}{i} p_{j-i}^{-j,-1}(x) + \frac{j-1}{i} (x-1) p_{j-i-1}^{-j+1,-1}(x) \right) = x p_{j-i}^{-j,0}(x) - p_{j-i}^{-j+2,-2}(x)$$

or

$$2 \left(\frac{j}{i} p_{j-i}^{-j,-1}(x) + \frac{j-1}{i} p_{j-i-1}^{-j+1,-1}(x) \right) = (x+1) p_{j-i}^{-j,0}(x) - (x-1) p_{j-i}^{-j+2,-2}(x)$$

It is now straightforward to derive this identity by using the formulas (4.5.1) and (4.5.4) from [Sz]. $\square$

5. APPENDIX: JACOBI AND GEGENBAUER POLYNOMIALS

The *Jacobi* polynomials $P_n^{\alpha,\beta}$ are defined for non negative integers n and arbitrary (rational) numbers α and β as follows (see Chapter IV in [Sz]),

$$P_n^{\alpha,\beta}(x) = \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} (n+\alpha+\beta+1) \dots (n+\alpha+\beta+k) \times (\alpha+k+1) \dots (\alpha+n) \left(\frac{x-1}{2} \right)^k,$$

with the understanding that the general coefficient

$$\binom{n}{k} (n+\alpha+\beta+1) \dots (n+\alpha+\beta+k) (\alpha+k+1) \dots (\alpha+n)$$

is equal to $(\alpha+1) \dots (\alpha+n)$ if $k=0$, and equal to $(n+\alpha+\beta+1) \dots (2n+\alpha+\beta)$ if $k=n$. The Jacobi polynomials can also be represented as

$$P_n^{\alpha,\beta}(x) = \frac{(\alpha+1) \dots (n+\alpha)}{n!} {}_2F_1 \left(-n, n+\alpha+\beta+1; \alpha+1; \frac{1-x}{2} \right),$$

where ${}_2F_1$ is the hypergeometric function of Gauss. If $\alpha = \beta$ the normalized Jacobi polynomials

$$\frac{\Gamma(\alpha+1)}{\Gamma(2\alpha+1)} \frac{\Gamma(n+2\alpha+1)}{\Gamma(n+\alpha+1)} P_n^{\alpha,\alpha}(x)$$

are called *ultraspherical* polynomials or *Gegenbauer's* polynomials. In what follows we shall call *ultraspherical Jacobi polynomials* to the “unnormalized” Gegenbauer's polynomials $P_n^{\alpha,\alpha}(x)$.

Let us consider

$$p_n^{\alpha,\beta}(x) = (x-1)^n P_n^{\alpha,\beta} \left(\frac{x+1}{x-1} \right),$$

it is clear that $p_n^{\alpha,\beta}(x)$ is again a polynomial. These polynomials can be expressed in terms of the hypergeometric function of Gauss as follows (see (4.22.1) in [Sz]),

$$p_n^{\alpha,\beta}(x) = \frac{(n + \alpha + \beta + 1) \dots (2n + \alpha + \beta)}{n!} {}_2F_1(-n, -n - \alpha; -2n - \alpha - \beta; x - 1).$$

5.1. Polynomial identities. It is well known that

$$(5.1) \quad P_n^{\alpha,\beta}(x) = (-1)^n P_n^{\beta,\alpha}(-x).$$

In particular, the ultraspherical Jacobi polynomials are even or odd according as n is even or odd. The ultraspherical Jacobi polynomials also satisfy the following identities (see (4.7.14) and (4.7.28) in [Sz]),

$$(5.2) \quad (n + 2\alpha)P_n^{\alpha,\alpha}(x) = 2(n + \alpha)P_n^{\alpha-1,\alpha-1}(x) + x(n + \alpha)P_{n-1}^{\alpha,\alpha}(x)$$

$$(5.3) \quad x(n + 2\alpha)P_n^{\alpha,\alpha}(x) = 2(n + 1)P_{n+1}^{\alpha-1,\alpha-1}(x) + (n + \alpha)P_{n-1}^{\alpha,\alpha}(x).$$

The difference between these two identities gives,

$$(5.4) \quad (1 - x)(n + 2\alpha)P_n^{\alpha,\alpha}(x) = -2(n + 1)P_{n+1}^{\alpha-1,\alpha-1}(x) + 2(n + \alpha)P_n^{\alpha-1,\alpha-1}(x) - (1 - x)(n + \alpha)P_{n-1}^{\alpha,\alpha}(x).$$

On the other hand, multiplying by x equation (5.2) and subtracting from it equation (5.3) yields the following identity,

$$(5.5) \quad 2(n + 1)P_{n+1}^{\alpha-1,\alpha-1}(x) = 2x(n + \alpha)P_n^{\alpha-1,\alpha-1}(x) + (x^2 - 1)(n + \alpha)P_{n-1}^{\alpha,\alpha}(x).$$

Lemma 5.1. *The ultraspherical Jacobi polynomials satisfy,*

$$\begin{aligned} 2 \sum_{k=0}^n \left(\frac{1-x}{2}\right)^{n-k} P_k^{\alpha+n-k,\alpha+n-k}(x) &= \frac{n+2\alpha}{n+\alpha} P_n^{\alpha,\alpha}(x) + P_{n-1}^{\alpha,\alpha}(x) \\ 2 \sum_{k=0}^n \left(\frac{-1-x}{2}\right)^{n-k} P_k^{\alpha+n-k,\alpha+n-k}(x) &= \frac{n+2\alpha}{n+\alpha} P_n^{\alpha,\alpha}(x) - P_{n-1}^{\alpha,\alpha}(x) \end{aligned}$$

Proof. The second identity follows immediately from the first one by using (5.1), then we shall prove the first identity by induction on n . It is clear that this identity holds for $n = 0$ and all α , hence if we assume that it holds for n and all α we have,

$$\begin{aligned} &2 \sum_{k=0}^{n+1} \left(\frac{1-x}{2}\right)^{n+1-k} P_k^{\alpha+n+1-k,\alpha+n+1-k}(x) \\ &= \frac{1-x}{2} 2 \sum_{k=0}^n \left(\frac{1-x}{2}\right)^{n-k} P_k^{\alpha+n+1-k,\alpha+n+1-k}(x) + 2P_{n+1}^{\alpha,\alpha}(x) \\ &= \frac{1-x}{2} \frac{n+2\alpha+2}{n+\alpha+1} P_n^{\alpha+1,\alpha+1}(x) + \frac{1-x}{2} P_{n-1}^{\alpha+1,\alpha+1}(x) + 2P_{n+1}^{\alpha,\alpha}(x) \\ &= -\frac{n+1}{n+\alpha+1} P_{n+1}^{\alpha,\alpha}(x) + P_n^{\alpha,\alpha}(x) + 2P_{n+1}^{\alpha,\alpha}(x) \\ &= \frac{n+1+2\alpha}{n+1+\alpha} P_{n+1}^{\alpha,\alpha}(x) + P_n^{\alpha,\alpha}(x), \end{aligned}$$

which completes the proof of the first identity. We point out that in the last but one equality we used the identity (5.4) with $\alpha + 1$ instead of α . $\square$

REFERENCES

- [BCT] A. Brega, L. Cagliero and J. Tirao, *The classifying ring of a classical rank one semisimple Lie group*, submitted.
- [C] Paul M. Cohn, *The similarity reduction of matrices over a skew field*, Math. Z. **132** (1973), 151–163.
- [CS] F. Chyzak and B. Salvy, *Noncommutative elimination in Ore algebras proves multivariate identities*, J. Symbolic Comput. **26** (1998), no. 2, 187–227.
- [GGRW] I. Gelfand, S. Gelfand, V. Retakh and R. Wilson, *Quasideterminants*, Adv. Math. **193** (2005), no. 1, 56–141.
- [KO] Koornwinder, T. and Onn, U., *LU factorizations, $q=0$ limits, and p -adic interpretations of some q -hypergeometric orthogonal polynomials*, Ramanujan J. **13** (2007), 365–387.
- [L] J. Lepowsky, *Algebraic results on representations of semisimple Lie groups*, Trans. Amer. Math. Soc. **176** (1973), 1–44.
- [Or] Oystein Ore, *Linear equations in noncommutative fields*, Ann. of Math. (2) **32** (1931), no. 3, 463–477.
- [Sz] G. Szegő, *Orthogonal polynomials*, AMS Colloquium Publications, Vol. XXIII (1959), New York.
- [T] J. Tirao, *On the centralizer of K in the universal enveloping algebra of $SO(n, 1)$ and $SU(n, 1)$* , Manuscripta math. **85** (1994), 119–139.

CIEM-FAMAF, UNIVERSIDAD NACIONAL DE CÓRDOBA
E-mail address: `brega@famaf.unc.edu.ar`

CIEM-FAMAF, UNIVERSIDAD NACIONAL DE CÓRDOBA
E-mail address: `cagliero@famaf.unc.edu.ar`